arXiv:solv-int/9512005v1 11 Dec 1995

On discrete 3-dimensional equations associated with the local Yang-Baxter relation

R.M. Kashaev*

Laboratoire de Physique Théorique ENSLAPP†
ENSLyon, 46 Allée d'Italie,
69007 Lyon, FRANCE
E-mail: `rkashaev@enslapp.ens-lyon.fr`

December 1995

Abstract

The local Yang-Baxter equation (YBE), introduced by Maillet and Nijhoff, is a proper generalization to 3 dimensions of the zero curvature relation. Recently, Korepanov has constructed an infinite set of integrable 3-dimensional lattice models, and has related them to solutions to the local YBE. The simplest Korepanov's model is related to the star-triangle relation in the Ising model. In this paper the corresponding discrete equation is derived. In the continuous limit it leads to a differential 3d equation, which is symmetric with respect to all permutations of the three coordinates. A similar analysis of the star-triangle transformation in electric networks leads to the discrete bilinear equation of Miwa, associated with the BKP hierarchy.

Some related operator solutions to the tetrahedron equation are also constructed.

ENSLAPP-L-569/95

---

*On leave of absence from St. Petersburg Branch of the Steklov Mathematical Institute, Fontanka 27, St. Petersburg 191011, RUSSIA

†URA 14-36 du CNRS, associée à l'E.N.S. de Lyon, au LAPP d'Annecy et à l'Universitè de Savoie

1 Introduction

The Yang-Baxter equation (YBE) [1, 2] plays the central role in the formalism of the quantum inverse scattering method of solving integrable 2d quantum systems [3]. The corresponding dynamical equations can be represented as a zero-curvature condition (Lax equation) for some auxiliary linear problem.

The notion of the local YBE has been introduced by Maillet and Nijhoff in [4, 5] as a proper 3-dimensional generalization of the zero-curvature condition. Korepanov in [6, 7] has pushed forward this idea by constructing an infinite set of discrete 3-dimensional integrable systems, and relating them to solutions to the local YBE, the simplest solution being equivalent to the star-triangle relation (STR) in the 2-dimensional Ising model.

In this letter first we consider the star-triangle transformation in electric networks, and show that, being interpreted as the local YBE, it leads to the integrable discrete bilinear equation, first discovered by Miwa in [8]. This is the content of Sect. 2. Then, in Sect. 3, in the same way we analyse the Korepanov's model associated with the STR in the Ising model. The corresponding integrable discrete equation in the continuous limit leads to the 3-dimensional non-linear differential equation, the symmetry group of which contains all permutations of three coordinates. In Sect. 4 some operator solutions to the tetrahedron equation, related with the abovementioned models, are constructed.

2 The star-triangle transformation in electric networks and Miwa's equation

The star-triangle transformation in electric networks is historically the first transformation of this kind known since the last century. It is given by the following formulae:

$$r_j r'_j = r'_1 r'_2 + r'_2 r'_3 + r'_3 r'_1 = r_1 r_2 r_3/(r_1 + r_2 + r_3), \quad j = 1,2,3, \tag{2.1}$$

where the resistances $\{r'_j\}$ and $\{r_j\}$ correspond to the star and triangle, respectively. There is a STR in the usual sense [9], underlying this transformation. Define a Gaussian exponent:

$$w(r|q) = \exp(-iq^2/2r), \quad q, r \in R. \tag{2.2}$$

Then, it is easy to check that the following STR holds:

$$\begin{aligned}&\int_{-\infty}^{+\infty} dq'' w(r'_3|q-q'')w(r'_2|q'')w(r'_1|q''-q')\\&= \sqrt{2\pi r'_1 r'_3/ir_2}\,w(r_1|q)w(r_2|q-q')w(r_3|q'),\end{aligned} \tag{2.3}$$

where the parameters $\{r'_j\}$, $\{r_j\}$ satisfy relations (2.1). The STR can be rewritten in the operator form as follows.

Let $C[\hat{p},\hat{q}]$ be the Heisenberg algebra generated by Hermitian operators $\hat{p}$ and $\hat{q}$, satisfying the Heisenberg permutation relation:

$$\hat{q}\hat{p} - \hat{p}\hat{q} = i. \tag{2.4}$$

Then, interpreting the variables q, q', q'' in the STR (2.3) as the eigenvalues of the operator $\hat{q}$, one can write the STR in the form:

$$w(-1/r'_3|\hat{p})w(r'_2|\hat{q})w(-1/r'_1|\hat{p}) = w(r_1|\hat{q})w(-1/r_2|\hat{p})w(r_3|\hat{q}), \tag{2.5}$$

where the function $w(r|q)$ is defined in (2.2). A particular choice of the parameters, $r'_i r_i = -1$, $r_1 = r_3 = -r_2 = \pm 1$, in (2.5) gives a constant solution to the YBE, which is discussed in [11] in the context of the quantum discrete sine-Gordon model.

To interpret (2.5) as a local YBE, introduce a linearly independent set of vectors in R^3, $\{e_1, e_2, e_3\}$, and the 3-dimensional lattice, generated by these vectors with integer coefficients:

$$M = \{n = n_1e_1 + n_2e_2 + n_3e_3 \mid n_1, n_2, n_3 \in Z\} \tag{2.6}$$

Let a_1, a_2, a_3 be the functions on M:

$$a_j \colon M \ni n \mapsto a_j(n) \in C, \quad j = 1, 2, 3.$$

The form of the identity (2.5) suggests the following parametrization:

$$r'_1 = -1/a_1(n),\ r'_2 = -a_2(n - e_1 - e_3),\ r'_3 = -1/a_3(n),$$

$$r_1 = a_1(n - e_2 - e_3),\ r_2 = 1/a_2(n),\ r_3 = a_3(n - e_1 - e_2).$$

Substituting these expressions into (2.5), we obtain the local YBE as is defined in [4, 5]:

$$\begin{aligned}&w(a_3(n)|\hat{p})w(-a_2(n-e_1-e_3)|\hat{q})w(a_1(n)|\hat{p})\\&= w(a_1(n-e_2-e_3)|\hat{q})w(-a_2(n)|\hat{p})w(a_3(n-e_1-e_2)|\hat{q}).\end{aligned} \tag{2.7}$$

The star-triangle transformation formulae (2.1) acquire now the form of difference equations:

$$\frac{a_j(n-e_k-e_l)}{a_j(n)} = -\frac{1+a_2(n-e_1-e_3)(a_1(n)+a_3(n))}{a_1(n)a_3(n)}, \ \{j,k,l\} = \{1,2,3\}. \tag{2.8}$$

Theorem 1 *The substitution*

$$a_j(n) = \alpha_j^{-1}\frac{\tau(n+e_k)\tau(n+e_l)}{\tau(n+e_1+e_2+e_3)\tau(n-e_j)}, \quad \{j,k,l\} = \{1,2,3\},$$

where α_j are some numerical coefficients, reduces system of equations (2.8) to the only equation on the function $\tau(n)$:

$$\sum_{j=1}^{4}\alpha_j\tau(n+e_j)\tau(n-e_j) = 0, \quad e_4 = -e_1-e_2-e_3, \quad \alpha_4 = \alpha_1\alpha_2\alpha_3. \tag{2.9}$$

The proof is through the straightforward substitution. Miwa in [8] has shown that equation (2.9) in the suitable continuous limit reduces to the BKP equation, while it itself is equivalent to the whole BKP hierarchy. The limit $\epsilon \to 0$, where $\alpha_j = \beta_j\epsilon$, β_j being fixed, reduces equation (2.9) to the Hirota's equation of the discrete Toda system [12]:

$$\sum_{j=1}^{3}\beta_j\tau(n+e_j)\tau(n-e_j) = 0. \tag{2.10}$$

3 A difference equation associated with the STR in the Ising model

In this section we make a similar analysis to that of sect. 2 of another STR, associated with the Ising model. The corresponding dynamical system has been introduced and studied by Korepanov in [6, 7], where he has proved that this system has enough set of conserved quantities to be integrable, and described how to construct solutions of the finite gap type.

Define the Bolzmann weights of the Ising model as a function W:

$$W\colon C\times\{1,-1\} \ni (x,\sigma) \mapsto W(x|\sigma) = 1+x+\sigma-x\sigma \in C, \tag{3.1}$$

The STR has the form of the following identity:

$$\begin{aligned}\sum_{\sigma''=\pm1} & W(x'_3|\sigma\sigma'')W(x'_2|\sigma'')W(x'_1|\sigma''\sigma') \\ &= (1+x'_1x'_2x'_3)W(x_1|\sigma)W(x_2|\sigma\sigma')W(x_3|\sigma'),\end{aligned} \tag{3.2}$$

where

$$x_jx_k = (x'_jx'_k + x'_l)/(1+x'_1x'_2x'_3), \quad \{j,k,l\} = \{1,2,3\}. \tag{3.3}$$

To rewrite STR (3.2) in a matrix form, first define the algebra $C[Z,X]$ of 2-by-2 matrices, generated by matrices Z and X, satisfying the relations:

$$Z^2 = X^2 = 1, \quad ZX = -XZ.$$

With these definitions, the STR (3.2) can be rewritten in the form (see also [10]):

$$\begin{aligned}W((1-x'_3)/(1+x'_3)|X)W(x'_2|Z)W((1-x'_1)/(1+x'_1)|X) \\ = RW(x_1|Z)W((1-x_2)/(1+x_2)|X)W(x_3|Z),\end{aligned} \tag{3.4}$$

where

$$R = \frac{(1+x_2)(1+x'_1x'_2x'_3)}{(1+x'_1)(1+x'_3)}.$$

Next, introduce the new parameters through the formulae:

$$4\xi_j = (x_j - 1/x_j)^2, \quad 4/\xi'_j = (x'_j - 1/x'_j)^2, \quad j = 1,2,3. \tag{3.5}$$

Consider three complex valued functions $b_j(n)$, $j = 1,2,3$, on the 3d lattice M defined in (2.6), and identify their values with the above defined parameters as follows:

$$\begin{gathered}\xi_1 = 1/b_1(n), \quad \xi_2 = b_2(n-e_1-e_3), \quad \xi_3 = 1/b_3(n), \\ \xi'_1 = 1/b_1(n-e_2-e_3), \quad \xi'_2 = b_2(n), \quad \xi'_3 = 1/b_3(n-e_1-e_2),\end{gathered} \tag{3.6}$$

Operator identity (3.4) now takes the form of the local YBE.

Theorem 2 *Under the substitution*

$$b_j(n) = -\frac{f(n+e_k)f(n+e_l)}{f(n+e_1+e_2+e_3)f(n-e_j)}, \quad \{j,k,l\} = \{1,2,3\},$$

the star-triangle transformation formulae (3.3) together with (3.5) and (3.6) imply the following equation on the function $f(n)$:

$$2\sum_{j=1}^{4}(f(n+e_j)f(n-e_j))^2+4\prod_{j=1}^{4}f(n+e_j)+4\prod_{j=1}^{4}f(n-e_j)$$
$$=\left(\sum_{j=1}^{4}f(n+e_j)f(n-e_j)\right)^2, \quad e_4=-e_1-e_2-e_3. \qquad (3.7)$$

To prove the theorem one has to exclude the dependent variables by calculating the corresponding resultants, which in intermediate steps are cumbersome. In these calculations the Maple computer system has been used.

Consider now a particular continuous limit of the equation (3.7). Let ϵ be the lattice spacing. Introduce the field

$$\phi(\epsilon n)=3\log(f(n)), \quad n\in M. \qquad (3.8)$$

Theorem 3 *In the limit $\epsilon\to 0$ with $x=(x_1,x_2,x_3)=\epsilon n\in R^3$ fixed, the equation (3.7) for the function (3.8) is reduced to the following differential equation:*

$$6\Delta_2\phi\sum_{j=1}^{4}(\partial_j^2\phi)^2=2(\Delta_3\phi)^2+(\Delta_2\phi)^3+8\sum_{j=1}^{4}(\partial_j^2\phi)^3, \qquad (3.9)$$

where $\phi=\phi(x)$, $x=(x_1,x_2,x_3)$,

$$\Delta_a=\sum_{j=1}^{4}\partial_j^a, \quad a=2,3,$$

$$\partial_j=\partial/\partial x_j, \quad j=1,2,3; \quad \partial_4=-\partial_1-\partial_2-\partial_3.$$

To prove the theorem one has to expand both sides of (3.7) in power series of ϵ up to the order of ϵ^6.

Thus, we have obtained a differential equation which should be integrable as a continuous limit of the discrete integrable equation, and which is symmetric with respect to any permutation of the three coordinates in R^3.

4 Solutions to the tetrahedron equation

The quantum theory of discrete systems, associated with the local YBE, should be described somehow by solutions to the tetrahedron equation (TE). Here we show how one can construct certain operator solutions to the TE using the star-triangle transformations considered in sections 2 and 3.

First, consider the case associated with electric networks. Let R_+ be the set of positive real numbers:

$$R_+ = \{r \in R \mid r > 0\},$$

and ι be the standard involution in it:

$$\iota\colon R_+ \to R_+, \quad \iota^*(r) = 1/r, \tag{4.1}$$

where the variable r is interpreted as a coordinate function on R_+. Define the following rational involution in R_+^3:

$$\varphi\colon R_+^3 \to R_+^3, \quad \varphi^*(r_j) = r_j \frac{r_1 + r_2 + r_3}{r_1 r_2 r_3}, \quad j = 1, 2, 3, \tag{4.2}$$

where r_1, r_2, r_3 are the coordinate functions of the corresponding multiples in R_+^3. Note, that φ commutes with the action of the permutation group on R_+^3. Formulas (4.2) are derived in fact from the star-triangle transformation (2.1) through the following identification:

$$r'_j = 1/\varphi^*(r_j), \quad j = 1, 2, 3.$$

Define now one more involution of R_+^3 as the following composition:

$$\Phi = \iota_2 \circ \varphi \circ \iota_2\colon R_+^3 \to R_+^3, \tag{4.3}$$

where $\iota_2 = \mathrm{id} \times \iota \times \mathrm{id}$. Let Φ_{ijk}, $1 \le i < j < k \le 6$, be the involutions of R_+^6, which act as Φ on the subspace $R_+^3 \subset R_+^6$, specified by the i-th, j-th and k-th R_+-multipliers, and identically, on the others.

Theorem 4 *The following TE holds in R_+^6:*

$$\Phi_{123} \circ \Phi_{145} \circ \Phi_{246} \circ \Phi_{356} = \Phi_{356} \circ \Phi_{246} \circ \Phi_{145} \circ \Phi_{123}. \tag{4.4}$$

The proof is the straightforward verification. The identity (4.4) implies the operator identity

$$\Phi_{123}^* \Phi_{145}^* \Phi_{246}^* \Phi_{356}^* = \Phi_{356}^* \Phi_{246}^* \Phi_{145}^* \Phi_{123}^* \tag{4.5}$$

in the linear space $\mathcal{F}(R_+^6)$ of complex-valued functions on R_+^6. Thus, we have obtained an operator solution to the TE. Solutions of such kind, called solutions to the "functional" TE, were considered also by Korepanov [13].

Let us comment on the limit, corresponding to Hirota's equation (2.10). For $\lambda \in R_+$ define the "rescaling" mapping:

$$g_\lambda : R_+^3 \to R_+^3, \quad g_\lambda^*(r_j) = \lambda r_j, \quad j = 1, 2, 3,$$

and consider the following composition:

$$\Phi_\lambda = g_{1/\lambda} \circ \Phi \circ g_\lambda : R_+^3 \to R_+^3,$$

where Φ is defined in (4.3), (4.2), (4.1). Clearly, Φ_λ satisfies the TE (4.4) for any $\lambda \neq 0$. The limit $\lambda \to 0$, however, is non-singular: $\lim_{\lambda\to 0} \Phi_\lambda = \Phi_0$, and the mapping Φ_0, associated with the Hirota's equation (2.10), satisfies (4.4) as well.

Consider now the star-triangle transformation (3.3), corresponding to the Ising model. Let I be the open unit interval in R:

$$I = \{x \in R \mid 0 < x < 1\},$$

and $\jmath$ be the following involution in I:

$$\jmath : I \to I, \quad \jmath^*(x) = (1 - x)/(1 + x), \tag{4.6}$$

where x is considered as a coordinate function on I. Define the following involution in I^3:

$$\psi : I^3 \to I^3,$$

$$\psi^*(x_j) = \sqrt{\frac{(1 - x_k x_l)^2 - x_j^2(x_k - x_l)^2}{(1 + x_k x_l)^2 - x_j^2(x_k + x_l)^2}}, \quad \{j, k, l\} = \{1, 2, 3\}. \tag{4.7}$$

This mapping is related with (3.3) through the following identification:

$$\psi^*(x_j) = (1 - x_j')/(1 + x_j').$$

The mapping

$$\Psi = \jmath_2 \circ \psi \circ \jmath_2 : I^3 \to I^3, \tag{4.8}$$

should satisfy the TE identity (4.4) with Φ's replaced by Ψ's and considered in I^6. We have verified this identity numerically, and claim it as a conjecture. The corresponding mapping

$$\Psi^* : \mathcal{F}(I^3) \to \mathcal{F}(I^3)$$

can be interpreted as a linear operator in the functional space $\mathcal{F}(I^3)$.

5 Summary

Integrable discrete equations in three dimensions can be obtained through solutions to the local YBE. Particularly, the star-triangle transformation in electric networks (2.1) can be associated with a solution to the local YBE (2.7), which leads to the integrable bilinear Miwa's equation (2.9). For the Korepanov's dynamical model, related with the star-triangle transformation in the Ising model, in this way we obtain another discrete equation (3.7). The continuous limit of the latter is described by a three dimensional differential equation (3.9), which is symmetric with respect to any permutation of coordinate axes.

The operator solutions (4.3), (4.2), (4.1), and (4.8), (4.7), (4.6) to the TE equation (4.4), connected with these dynamical models, may help to quantize the latter. Note, that the quantum version of Hirota's equation (2.10) has been constructed in [14] in the context of the quantum inverse scattering method.

6 Acknowlegements

The author is grateful to L.D. Faddeev and I.G. Korepanov for the stimulating conversations, and to J.M. Maillet for the discussion of the local YBE. The work is supported by the Programme TEMPRA-Europe de l'Est from the Région Rhône-Alpes.

References

[1] C.N. Yang, *Some exact results for the many-body problem in one dimension with repulsive delta-function interaction*, Phys. Rev. Lett. 1967, Vol. 19, P. 1312-1314

[2] R.J. Baxter, *Partition function of the eight-vertex lattice model*, Ann. Phys. 1972, Vol. 70, P. 193-228

[3] L.D. Faddeev, *Quantum completely integrable models in field theory*, Sov. Sci. Rev. C**1** (1980) 107-155

[4] J.M. Maillet, F.W. Nijhoff, *Integrability for multidimensional lattice models*, Phys. Lett. B224 (1989) 389-396

[5] J.M. Maillet, *Integrable systems and gauge theories*, Nucl. Phys. (Proc. Suppl.) 18B (1990) 212-241

[6] I.G. Korepanov, *A dynamical system connected with inhomogeneous 6-vertex model*, Zap. Nauch. Semin. POMI 215 (1994) 178-196

[7] I.G. Korepanov, *Algebraic integrable dynamical systems, 2+1-dimensional models in wholly discrete space-time, and inhomogeneous models in 2-dimensional statistical physics*, solv-int/9506003

[8] T. Miwa, *On Hirota's Difference Equations*, Proc. Japan Acad., 58, Ser. A (1982) 9-12

[9] R.J. Baxter, *Exactly solved models in statistical mechanics*, N.Y., Academic Press, 1982

[10] V.B. Matveev, A.O. Smirnov, *Some comments on the solvable chiral Potts model*, Lett. Math. Phys. 19 (1990) 179-185

[11] L.D. Faddeev, *Current-like variables in massive and massless integrable models*, Lectures delivered at the International School of Physics "Enrico Fermi", held in Villa Monastero, Varenna, Italy, 1994.

[12] R. Hirota, *Discrete analogue of a generalized Toda equation*, J. Phys. Soc. Jpn., Vol. 50, No. 11, (1981) 3785-3791

[13] I.G. Korepanov, private communication, 1995

[14] R.M. Kashaev, N.Yu. Reshetikhin, *Affine Toda field theory as a 3-dimensional integrable system*, preprint ENSLAPP-L-548/95